

Emission estimates of selected volatile organic compounds from tropical savanna burning in northern Australia

T. Shirai,¹ D. R. Blake,² S. Meinardi,² F. S. Rowland,² J. Russell-Smith,³ A. Edwards,³ Y. Kondo,⁴ M. Koike,⁵ K. Kita,⁶ T. Machida,⁷ N. Takegawa,⁸ N. Nishi,⁹ S. Kawakami,¹ and T. Ogawa¹

Received 17 May 2001; revised 22 December 2001; accepted 8 February 2002; published 24 January 2003.

[1] Here we present measurements of a range of carbon-based compounds: carbon dioxide (CO₂), carbon monoxide (CO), methane (CH₄), nonmethane hydrocarbons (NMHCs), methyl halides, and dimethyl sulfide (DMS) emitted by Australian savanna fires studied as part of the Biomass Burning and Lightning Experiment (BIBLE) phase B aircraft campaign, which took place during the local late dry season (28 August to 13 September 1999). Significant enhancements of short-lived NMHCs were observed in the boundary layer (BL) over the region of intensive fires and indicate recent emissions for which the mean transport time was estimated to be about 9 hours. Emission ratios relative to CO were determined for 20 NMHCs, 3 methyl halides, DMS, and CH₄ based on the BL enhancements in the source region. Tight correlations with CO were obtained for most of those compounds, indicating the homogeneity of the local savanna source. The emission ratios were in good agreement with some previous measurements of savanna fires for stable compounds but indicated the decay of emission ratios during transport for several reactive compounds. Based on the observed emission ratios, emission factors were derived and compared to previous studies. While emission factors (g species/kg dry mole) of CO₂ varied little according to the vegetation types, those of CO and NMHCs varied significantly. Higher combustion efficiency and a lower emission factor for methane in this study, compared to forest fires, agreed well with results for savanna fires in other tropical regions. The amount of biomass burned was estimated by modeling methods using available satellite data, and showed that 1999 was an above average year for savanna burning. The gross emissions of the trace gases from Australian savanna fires were estimated. *INDEX TERMS*: 0365 Atmospheric Composition and Structure: Troposphere—composition and chemistry; 0368 Atmospheric Composition and Structure: Troposphere—constituent transport and chemistry; 0322 Atmospheric Composition and Structure: Constituent sources and sinks

Citation: Shirai, T., et al., Emission estimates of selected volatile organic compounds from tropical savanna burning in northern Australia, *J. Geophys. Res.*, 108(D3), 8406, doi:10.1029/2001JD000841, 2003.

¹National Space Development Agency of Japan, Earth Observation Research Center, Tokyo, Japan.

²Department of Chemistry, University of California, Irvine, California, USA.

³Tropical Savannas Cooperative Research Centre, Bushfires Council of the Northern Territory, Northern Territory, Australia.

⁴Research Center for Advanced Science and Technology, University of Tokyo, Tokyo, Japan.

⁵Department of Earth and Planetary Sciences, University of Tokyo, Tokyo, Japan.

⁶Department of Environmental Sciences Faculty of Science, Ibaraki University, Ibaraki, Japan.

⁷National Institute for Environmental Studies, Tsukuba, Japan.

⁸Solar-Terrestrial Environment Laboratory, Nagoya University, Aichi, Japan.

⁹Department of Earth and Planetary Sciences, Kyoto University, Kyoto, Japan.

1. Introduction

[2] Biomass burning is a globally significant source of important atmospheric trace gases such as CO, CO₂, CH₄, NMHCs, nitrogen oxides (NO_x = NO + NO₂), halogenated and oxygenated compounds, and aerosol particles, which together affect atmospheric chemistry and climate on regional and global scales [Crutzen and Andreae, 1990; Granier et al., 1999; Yokelson et al., 1999a].

[3] The photochemical oxidation of CO and volatile organic compounds (VOCs) in the presence of NO_x is a major source of tropospheric ozone (O₃). Tropospheric O₃ is an important greenhouse gas as well as a major precursor of the hydroxyl radical (OH) which plays a key role in tropospheric chemistry. Biomass burning is also a major source of the methyl halides; methyl chloride (CH₃Cl), methyl bromide (CH₃Br), and a minor source of methyl iodide (CH₃I). The methyl halides have come to the forefront in the last decade due to their potential for delivering ozone-depleting halogens

to the stratosphere [Andreae et al., 1996; Blake et al., 1996a; World Meteorological Organization (WMO), 1999].

[4] About 90% of global biomass burning occurs in the tropics and extensive field studies on the atmospheric effects of biomass burning have been conducted during the last decade [Hao and Liu, 1994; Andreae et al., 1994; Folkins et al., 1995; Browell et al., 1996; Connors et al., 1996; Talbot et al., 1996; Jonquière et al., 1998; Blake et al., 1999; Gregory et al., 1999; Singh et al., 2000]. In northern Australia, biomass burning has occurred regularly in the dry season for many thousands of years. These fires called “bushfires” occur in a tropical eucalyptus savanna dominated by eucalyptus trees, small bushes and grasses [Lacey et al., 1982] and are mainly anthropogenic in origin. Although biomass burning in Australia is a significant source of gases with regional and global influences on the troposphere, few studies have reported emissions from Australian savanna fires, as compared with Africa and tropical America [Hurst et al., 1994a, 1994b]. In this paper, we estimate the annual gross emissions of several important atmospheric compounds (e.g., CO₂, CO, CH₄, NMHCs, methyl halides) from savanna burning in Australia, based on the trace gas measurements in smoke emitted from savanna burning in the late dry season.

2. Experiment

[5] Observations of atmospheric trace gases were made utilizing a Gulf stream-II (G-II) aircraft as part of BIBLE (Biomass Burning and Lightning Experiment) phase B sponsored by NASDA (National Space Development Agency of Japan) during the local late dry season (28 August to 13 September 1999). The campaign consisted of 1 test flight over Japan, 6 ferry flights between Japan and Australia, and 6 local flights over northern Australia to survey trace gas emissions from savanna burning. During the entire campaign, more than 600 whole air samples were collected aboard the G-II in 2-L stainless steel canisters. In situ measurements of O₃, CO, CO₂, NO, NO_y (= NO + NO₂ + NO₃ + N₂O₅ + PAN + HNO₂ + HNO₃ + HNO₄ + other nitrates), aerosols, condensation nuclei, soot particles, and UV flux were also made by other collaborating groups.

[6] The whole air samples were returned to a laboratory at the University of California, Irvine (UCI) and were analyzed by multicolumn gas chromatography. Two flame ionization detectors (FIDs) and two electron capture detectors (ECDs) in combination with four gas chromatographic (GC) columns were optimized for the determination of selected compounds. In addition, the system employed a bench-top quadrupole mass spectrometer (MS) to help positively identify unknown components. The FIDs were used to quantify C₂-C₈ nonmethane hydrocarbons (NMHCs) while the ECDs and MS detected halocarbons and alkyl nitrates. Most quantified gases were detected on more than one of the column/detector combinations, which provide an additional check on proper identification and quantification. Fifty-three compounds have been archived, including 24 halocarbons, 23 nonmethane hydrocarbons (NMHCs), 5 alkyl nitrates and dimethyl sulfide (DMS). The accuracy of the measurements ranges from 5–10%. The limit of detection for the NMHCs was 3 parts per trillion by volume (pptv) and the measurement precisions

range from 1–5%. The accuracy, precision and the limit of detection vary for each compound and are presented in the overview paper of the BIBLE campaign [Kondo et al., 2002]. Further details about sampling procedures are given by Blake et al. [1992, 1994], and the experimental apparatus is described by Blake et al. [1996a], Sive [1998], and Colman et al. [2001]. CH₄ was analyzed by a separate GC/FID system with precision of 0.1%.

[7] During BIBLE-B, fast response in situ measurements of CO₂ and CO were made using a non-dispersive infrared analyzer (NDIR) and a resonance fluorescence technique, respectively. Details of the measurement techniques are described elsewhere [Machida et al., 2002; Takegawa et al., 2001]. Because measurement frequency of and sampling duration varied by species [Kondo et al., 2002], the data were averaged over a common time interval before data analyses. The merged data used in this study were generated based on the sampling time of the whole air samples, which had the lowest sampling frequency of all the measurements reported in this paper.

3. Results and Discussion

[8] During the late dry season when BIBLE-B was conducted, active bushfires were a daily occurrence in northern Australia, as shown by hot spot distributions in NOAA/AVHRR satellite imagery [Russell-Smith et al., 2003]. When the aircraft flew within the boundary layer (BL) below approximately 3 km over the source region north of the Northern Territory in Australia, it often passed through smoke plumes in which elevated mixing ratios of CO, CO₂, NO, and NO_y were observed. Typical CO values observed over the source region at altitudes <3 km were 160–200 ppbv while the values observed upwind of the source region were about 80 ppbv [Takegawa, 2003]. Figure 1 shows the time series of flight altitude and concentrations of selected trace gases observed during one of the local flights (flight 7 shown in Kondo et al., this issue) over northeastern Australia (10°–30°S, 120°–150°E). During the flight, CO values occasionally exceeded 1 ppmv and large enhancements of methyl halides were also observed in these plumes (data not shown). The mixing ratios of short-lived alkenes such as propene (C₃H₆) and butenes (1-C₄H₈, c-2-C₄H₈, and t-2-C₄H₈) observed in these plumes reached up to 2.8 ppbv, 390 pptv, 53 pptv, and 67 pptv, respectively, which indicate very rapid convection from active biomass fires [Blake et al., 1996a].

[9] Figure 2 illustrates the correlations of selected trace gases versus CO for the BL subset of two intensive flights which deployed from Darwin. The data set consists of 92 samples collected over Arnhemland, which was strongly influenced by biomass burning emissions. Forty-three samples were collected at 1.7 km altitude during a flight conducted on 2 September, and 49 samples were collected during a flight conducted on 4 September, with 31 samples collected at 1.7 km and 18 samples at 3.0 km. The two flight paths are shown in Figure 3 together with the AVHRR hot spots observed close to the flight deployments at similar times.

[10] As shown in Figure 2, most of the trace gases such as CO₂, CO, CH₄, NMHCs, and methyl halides that are known to be produced in fires were highly correlated with each other ($r^2 > 0.9$). These tight correlations indicate the identical




Figure 1. Time series of ethane, propene and carbon monoxide (CO) mixing ratios observed during local flight 7 over northern Australia on 4 September 1999. Thick solid line traces flight altitude.

source for those compounds, which apparently represents local savanna burning emissions. The urban tracer C_2Cl_4 showed no correlation with CO, indicating no contribution from urban sources (Figure 2). Aircraft measurements made in well-mixed downwind smoke made during this study should provide good integrated samples with which to evaluate the emissions of relatively stable compounds as indicated by the tight correlations shown in Figure 2.

[11] The possible contribution from oceanic sources to these samples was dismissed by the good correlations of the compounds that have both oceanic and biomass burning sources such as CH_3Cl , CH_3I , and DMS with CO. Observed DMS/CO was 0.006 pptv/ppbv ($r^2 = 0.79$). DMS is a precursor of sulfuric acid and particulate sulfate in the marine atmosphere, which affects the global radiation budget, rain acidity, etc [Charlson *et al.*, 1987; Warneck, 2000]. Although oceanic emissions greatly outweigh DMS emissions from fires on a global scale, the good DMS to CO correlation that was observed indicates that emission from fires form a considerable part of DMS emission to the atmosphere in the study region (S. Meinardi *et al.*, Dimethyl disulfide and dimethyl sulfide emissions from biomass burning, submitted to *Science*, 2002, hereinafter referred to as Meinardi *et al.*, submitted manuscript, 2002).

[12] Methyl halides are emitted from various sources, both natural and anthropogenic [Butler, 2000]. Along with high correlations with CO, significant correlations were observed for CH_3Cl versus CH_3Br ($r^2 = 0.91$), CH_3Cl versus CH_3I ($r^2 = 0.91$), and CH_3Br versus CH_3I ($r^2 = 0.82$). Assuming that each source has its unique emission ratio, the single regression line indicates a single dominant source, which is assumed to be local biomass burning in this case.

[13] To discriminate between various burning stages, we calculated the combustion efficiency of the fire using the modified combustion efficiency (MCE), indicator, defined as $\Delta CO_2 / (\Delta CO_2 + \Delta CO)$ by [Ward and Radke, 1993]. We derived a mean MCE value of 0.92 for the aircraft samples

collected for this study. The observed aircraft MCE is within the range of typical values for savanna fires reported previously [Bonsang *et al.*, 1991; Hurst *et al.*, 1994a, 1994b; Andreae *et al.*, 1996; Blake *et al.*, 1996a]. The MCE value of 0.93 was observed for the ground samples collected during flaming stage of fires while much lower MCE (=0.81) values were observed for those collected during the smoldering stage of fires (Meinardi *et al.*, submitted manuscript, 2002). The MCE for the aircraft samples was much closer to the MCE for the ground samples representing the flaming stage, suggesting a possible bias associated with aircraft samples that may better represent easily convected hot flaming emissions rather than cooler smoldering emissions [Ferek *et al.*, 1998].

4. Emission Ratios

[14] Biomass burning emissions are commonly characterized by determining the ratio of $\Delta Y / \Delta X$, where y is a compound either emitted by the fire or produced in the fire plume, x is a reference tracer; usually CO or CO_2 ; and Δ indicates the excess mixing ratio in the plume over the local background.

[15] CO_2 is often employed as the reference tracer for flaming products such as H_2O , NO, and SO_2 , because the CO_2 emission rate is much higher in the flaming stage than in the smoldering stage. On the other hand, because CO is primarily emitted during smoldering, it is a good reference for other products of incomplete combustion, such as CH_4 , NMHCs, and NH_3 [Lobert *et al.*, 1991; Yokelson *et al.*, 1996]. Because several compounds (including light unsaturated hydrocarbons) are found to be uncorrelated temporally with either CO or CO_2 [Yokelson *et al.*, 1996, 1997] in the process of combustion, we must note that the use of CO as a reference might underestimate calculated emission ratios for certain compounds. However, the very good correlation between CO_2 and CO (Figure 2) indicates that these aircraft samples represent well-mixed emissions of flaming and




Figure 2. Correlation plots for savanna fire emissions observed during BIBLE-B.

smoldering combustion. A possible bias arising from using either CO or CO₂ as a reference gas is considered to be small.

[16] The emission ratios relative to CO for selected trace gases determined in this airborne observation are presented

in Table 1. The subset of 23 compounds listed were chosen from the larger set of compounds quantified with adequate precision and accuracy in this study, omitting the compounds that showed insignificant emission ratios. These emission ratios are the slopes of the correlation plots shown




Figure 3. BIBLE-B flight paths on 2 and 4 September 1999. NOAA AVHRR hot spots are indicated by the circles and squares (solid circles: 2 September, 19:20 WST, 1999, and solid squares: 4 September, 18:37 WST, 1999). WST: Australian Western Standard Time; 8 hours ahead of GMT or UTC.

in Figure 2, thus avoiding the need to subtract “background” mixing ratios. Errors indicated in Table 1 are standard errors $= \sqrt{\left(\frac{\sum_{i=1}^n x_i^2 - n\bar{x}^2}{n(n-1)}\right)}$.

[17] Emission ratios previously observed for African and Brazilian biomass burning are also presented in Table 1. These are the results from the Transport and Atmospheric Chemistry Near the Equator-Atlantic (TRACE-A) and the Brazil Fire Assessment (BFA) studies for which the experimental procedures were common with this study [Blake *et al.*, 1996b].

[18] Some general characteristics of emissions from Australian savanna fires are shown by the magnitude of the trace gas emission ratios (Table 1). For aliphatic compounds, emissions from savanna fires decrease with increasing carbon number. Among the series with the same carbon number, alkene emissions are higher than for alkanes or alkynes. Among aromatics, the emissions of benzene and toluene are prominent, and are comparable in magnitude to the emission of several C₃-C₄ aliphatic compounds. The highest emitted NMHC was C₂H₄, which accounts for 40% of the total NMHC emissions measured. The observed ratios agree fairly well with previous observations of African savanna fires for less reactive compounds which were little affected by decomposition during transport [Bonsang *et al.*, 1991; Blake *et al.*, 1996a; Hao *et al.*, 1996].

[19] The methyl halide emission ratios were well within the range of previous measurements (Table 1). Considering that these emissions are mainly controlled by halogen concentrations in fuels along with fire intensity and combustion efficiency [Reinhardt and Ward, 1995], the variations of emission ratios observed in different tropical regions were smaller than expected based on the large variations of halogen concentrations among fuels [McKenzie *et al.*, 1996]. As a result, it appears that the emissions of these compounds depend more on combustion efficiency than on halogen content in the fuels.

[20] When comparing emission ratios, the effects of photochemical processes and dilution must be taken into account. Emission ratios change during transport due to the different reactivity between the compound of interest and

the reference compound [Mauzerall *et al.*, 1998; Goode *et al.*, 2000]. This is apparent when the results of TRACE-A and BFA are compared (Table 1). While the emission ratios for relatively stable compounds such as alkanes vary little, those for very reactive alkenes vary considerably as a result of differences in transit times. Higher emission ratios for reactive compounds shown in the BFA results indicate shorter transport time compared to the TRACE-A observations [Blake *et al.*, 1996b].

[21] The factors that affect emission ratios were evaluated by comparing the emission ratios based on our aircraft observations with those based on ground samples collected at fires. Forty five whole air samples were collected on the ground at two different northern Australian savanna sites during the BIBLE-B experiment. These measurements are described in detail by Meinardi *et al.* (submitted manuscript, 2002). Both the magnitude and the variation of emission ratios for most of the observed trace gases were much higher for the ground samples than for aircraft samples. Compared to the background concentrations, the observed CO₂ enhancement was 100 times higher in ground samples and 1% higher in aircraft samples. For CO, the observed enhancement was 10,000 times higher in ground samples and 15 times higher in aircraft samples, compared to background levels.

[22] The variation of emission ratios observed for individual samples were greater for ground samples compared to aircraft samples. The larger variation among ground samples are because these samples were collected from a single fire over a relatively short sampling period (several minutes), and thus representing emissions from specific fuels at specific burning stages. By contrast, aircraft samples typically contain a mixture of emissions from multiple fires at various burning stages, thus representing emissions from many fires, or an average for the region. The emission ratios of each single fire deviate from the average according to fuel compositions and burning processes. However, for most of the ground samples taken during BIBLE-B, CO/CO₂ ratios were close to the aircraft CO/CO₂ value, except some samples that showed remarkably larger CO/CO₂ ratios. Those samples exhibiting extreme CO/CO₂ mixing

Table 1. Emission Ratios Relative to CO Observed for Tropical Biomass Burning (Based on Airborne Observation)

Name	Formula	Australia (BIBLE-B, 1999)		Brazil ^a (BFA, 1994)		Brazil ^b (TRACE-A, 1992)		Africa ^b (TRACE-A, 1992)	
		Ratio, pptv/ppbv	R ²	Ratio, pptv/ppbv	R ²	Ratio, pptv/ppbv	R ²	Ratio, pptv/ppbv	R ²
Methyl bromide	CH ₃ Br	5.9 ± 0.3 (× 10 ⁻³)	0.83	7.4 ± 0.9 (× 10 ⁻³)	0.49	10.6 ± 0.8 (× 10 ⁻³)	0.88	6.0 ± 0.5 (× 10 ⁻³)	0.78
Methyl chloride	CH ₃ Cl	0.38 ± 0.01	0.89	0.31 ± 0.03	0.90	0.85 ± 0.06	0.88	0.57 ± 0.03	0.92
Methyl iodide	CH ₃ I	2.8 ± 0.1 (× 10 ⁻³)	0.82	3.6 ± 0.3 (× 10 ⁻³)	0.60	1.2 ± 0.2 (× 10 ⁻³)	0.57	1.2 ± 0.2 (× 10 ⁻³)	0.70
Ethane	C ₂ H ₆	2.8 ± 0.1	0.97	8.5 ± 0.4	0.84	8.3 ± 0.3	0.97	5.2 ± 0.2	0.96
Ethene	C ₂ H ₄	7.1 ± 0.2	0.93	17.1 ± 0.4	0.95	6.1 ± 0.4	0.87	8.3 ± 0.6	0.78
Ethylene	C ₂ H ₂	2.9 ± 0.1	0.96	4.6 ± 0.2	0.90	3.3 ± 0.1	0.97	4.5 ± 0.1	0.98
Propane	C ₃ H ₈	0.58 ± 0.01	0.97	1.6 ± 0.1	0.79	1.6 ± 0.1	0.95	0.97 ± 0.04	0.94
Propene	C ₃ H ₆	1.6 ± 0.1	0.86	4.9 ± 0.1	0.98	0.4 ± 0.1	0.62	1.0 ± 0.1	0.56
Isobutane	i-C ₄ H ₁₀	0.048 ± 0.001	0.96	0.067 ± 0.004	0.78	0.055 ± 0.010	0.55	0.043 ± 0.004	0.77
Propadiene	C ₃ H ₄	0.054 ± 0.002	0.91	0.56 ± 0.04	0.76	0.25 ± 0.02	0.93	0.29 ± 0.02	0.86
Normal-Butane	n-C ₄ H ₁₀	0.11 ± 0.002	0.97	0.21 ± 0.02	0.70	0.21 ± 0.02	0.80	0.16 ± 0.01	0.88
Trans-2 Butene	t-2-C ₄ H ₈	0.026 ± 0.003	0.52	0.26 ± 0.01	0.88	-	-	-	-
1-Butene	1-C ₄ H ₈	0.23 ± 0.01	0.84	0.75 ± 0.02	0.75	0.06 ± 0.01	0.52	0.12 ± 0.04	0.54
Cis-2-Butene	c-2-C ₄ H ₈	0.022 ± 0.002	0.60	0.20 ± 0.01	0.87	-	-	-	-
Iso-Pentane	i-C ₅ H ₁₂	6.5 ± 0.5 (× 10 ⁻³)	0.69	0.020 ± 0.002	0.57	-	-	-	-
Normal-Pentane	n-C ₅ H ₁₂	0.028 ± 0.001	0.97	0.043 ± 0.002	0.84	0.051 ± 0.005	0.80	0.053 ± 0.005	0.80
Hexane	C ₆ H ₁₄	0.042 ± 0.002	0.89	-	-	-	-	-	-
Benzene	C ₆ H ₆	0.86 ± 0.02	0.97	1.7 ± 0.03	0.97	1.29 ± 0.05	0.97	1.30 ± 0.04	0.96
Heptane	C ₇ H ₁₆	0.017 ± 0.001	0.83	-	-	-	-	-	-
Toluene	C ₇ H ₈	0.28 ± 0.01	0.93	0.72 ± 0.02	0.72	0.57 ± 0.03	0.93	0.49 ± 0.03	0.85
Meta-Xylene	m-C ₈ H ₁₀	0.034 ± 0.002	0.85	-	-	0.042 ± 0.004	0.83	0.026 ± 0.007	0.52
Para-Xylene	p-C ₈ H ₁₀	9.8 ± 0.5 (× 10 ⁻³)	0.81	-	-	0.023 ± 0.003	0.75	0.014 ± 0.004	0.59
Ortho-Xylene	o-C ₈ H ₁₀	0.024 ± 0.001	0.90	-	-	0.027 ± 0.002	0.87	0.014 ± 0.004	0.53
Methane	CH ₄	0.044 ± 0.003 v/v	0.78	0.107 ± 0.004 v/v	0.86	0.094 ± 0.013 v/v	0.70	0.078 ± 0.004 v/v	0.91
CO/CO ₂		0.086 ± 0.002 v/v	0.97	(0.035) v/v	0.44	0.037 ± 0.003 v/v	0.87	0.062 ± 0.002 v/v	0.96

^aFrom Blake *et al.* [1996b].^bFrom Blake *et al.* [1996a].

v/v = ratio by volume.

These values are lower limits for reactive species; see text.

Table 2. Comparison of Emission Ratios Relative to CO in Aircraft Samples and Ground Samples^a

Compounds	BIBLE Aircraft E. Ratio, pptv/ppbv	BIBLE Ground E. Ratio, pptv/ppbv	Grd/Air Ratio (ER/ER)
<i>Lifetime Range in Months</i>			
C ₂ H ₆	2.8 ± 0.1	2.6 ± 0.5	0.93
<i>Lifetime Range in Weeks</i>			
C ₂ H ₂	2.9 ± 0.1	3.0 ± 0.2	1.03
C ₃ H ₈	0.58 ± 0.01	0.84 ± 0.12	1.45
C ₆ H ₆	0.86 ± 0.02	0.75 ± 0.03	0.87
<i>Lifetime Range in Days</i>			
n-C ₄ H ₁₀	0.11 ± 0.002	0.15 ± 0.02	1.36
i-C ₄ H ₁₀	0.048 ± 0.001	0.051 ± 0.007	1.06
n-C ₅ H ₁₂	0.028 ± 0.001	0.037 ± 0.006	1.32
C ₆ H ₁₄	0.042 ± 0.002	0.024 ± 0.003	0.57
C ₇ H ₈	0.28 ± 0.01	0.39 ± 0.04	1.39
<i>Lifetime Range in Hours</i>			
C ₂ H ₄	7.1 ± 0.2	8.2 ± 0.3	1.15
C ₃ H ₆	1.6 ± 0.1	2.3 ± 0.2	1.44
1-C ₄ H ₈	0.24 ± 0.01	0.39 ± 0.02	1.63
c-2-C ₄ H ₈	0.022 ± 0.002	0.084 ± 0.012	3.82
t-2-C ₄ H ₈	0.026 ± 0.003	0.11 ± 0.02	4.23

^aSeveral samples of extremely high CO/CO₂ ratios were excluded from the average (see text).

ratios were excluded from the comparison because they are biased to represent fires with extremely low combustion efficiency. Table 2 shows the emission ratios of selected NMHCs based on both aircraft samples and ground samples. Typical lifetimes for the conditions encountered during BIBLE-B for each compound are also shown in order to monitor the effect of destruction during transport, assuming the ground sample emission ratios represent initial values. Relative differences between ground and airborne ratios stay within ±60% for compounds with lifetimes longer than several hours, while up to 400% differences were observed for butene isomers with lifetimes of 2–3 hours (Table 2). In the following sections, only longer-lived compounds are taken into account when comparing emission ratios with previous studies.

[23] Using Y/CO (emission ratios of selected NMHCs) for ground samples and aircraft samples and the respective NMHC and CO lifetimes τ , the averaged air mass age from aircraft samples was calculated using $\ln r \cdot \tau$, ($r =$

ER(ground)/ER(aircraft), $\tau = k(\text{NMHC} + \text{OH})[\text{OH}]$), assuming the lifetime of CO is greater than the transport time. Lifetimes were calculated based on OH oxidation as the dominant sink process with an estimated mean [OH] value of 2.7×10^6 molecules cm⁻³ for the altitude range 0–4 km [Gregory *et al.*, 1996]. It should be noted that a lower OH concentration is expected in the biomass burning plume, which is influenced by high levels of CO and hydrocarbons as well as low J(O¹D) unless NO levels are high. However, in the plume we encountered during the BIBLE-B flights 6 and 7, when average NO levels were as high as 160 pptv (maximum 53 ppbv), it might well be that the reaction: HO₂ + NO → OH + NO₂ worked as a buffer and OH concentrations did not drop significantly [Ridley and Atlas, 1999]. In this estimation, emission ratios for C₂H₆, C₂H₄, C₂H₂, C₃H₈, C₃H₆, n-C₄H₁₀, 1-C₄H₈, C₆H₆, and C₇H₈ were used. The 2-butene isomers were excluded to avoid overestimation of air mass age by ignoring O₃ oxidation as a sink. The reaction contributes comparably to OH reaction for those internal olefins that react rapidly with O₃.

[24] The mean transport time of sampled air was estimated to be 0.4 days (8 hours) as an average of $\ln(Y/\text{CO}) \cdot \tau$ calculated for the series of compounds listed in the Table 2. This estimate is consistent with the significant enhancements of short-lived alkenes whose lifetimes are less than 0.2 days at the low altitudes at which the plume was sampled. However, for a more precise estimation, the effects of dilution and entrainment on NMHC concentrations by mixing with circumambient air during transport must be considered for aircraft samples, which remains to be an issue in the future.

[25] In Table 3, previous emission ratios from Australian savanna biomass burning (derived from the ratios between excess mixing ratios $\Delta Y/\Delta X$), are shown with our airborne values converted into the same units. The previous experiments were conducted also during the late dry season over the same region, using a light aircraft flying at 50–700 m above the fires [Hurst *et al.*, 1994a, 1994b]. While Hurst *et al.* [1994a, 1994b] reported multiple measurements made using different instruments, the values presented in the Table are the results of flask sampling and GC analysis only, which was a similar technique to ours. For CO, and CH₄, the BIBLE-B emission ratios were in generally good agreement with the previous values for Australian savannas. This good

Table 3. Enhancement Ratios Relative to CO for Australian Savanna Fires (Based on Airborne Observation)^a

	This Study		Hurst [1994a]		Hurst [1994b]	
	Ratio (pptv/ppbv)	R ²	Ratio (pptv/ppbv)	R ²	Ratio (pptv/ppbv)	R ²
CO/CO ₂	0.089 ± 0.002 v/v	0.97	0.090 ± 0.026 v/v	0.86	0.058 ± 0.020 v/v	0.93
CH ₄	0.043 ± 0.003 v/v	0.76	0.045 ± 0.009 v/v	0.95	0.068 ± 0.040 v/v	0.91
C ₂ H ₂	3.0 ± 0.1	0.97	0.72 ± 0.37	0.74	2.5 ± 1.1	0.98
C ₂ H ₄	7.5 ± 0.3	0.89	-	-	11 ± 5	0.97
C ₂ H ₆	2.7 ± 0.1	0.97	-	-	4.8 ± 1.8	0.96
C ₃ H ₆	1.6 ± 0.7	0.85	-	-	2.3 ± 1.0	0.98
C ₃ H ₈	0.57 ± 0.01	0.97	-	-	0.77 ± 0.19	0.98
n-C ₄ H ₁₀	0.11 ± 0.002	0.97	-	-	0.21 ± 0.04	0.99
i-C ₃ H ₁₂	6.7 ± 0.3 (×10 ⁻³)	0.80	-	-	0.25 ± 0.10	0.98
C ₆ H ₆	0.85 ± 0.01	0.97	0.19 ± 0.15	0.68	0.42 ± 0.24	0.98

^aThese values are lower limits for reactive species. See text.

Table 4. Emission Factors Observed for Various Vegetation Types (g Species/kg dm)

	Australian Savanna			Brazilian Cerrado ^b	Global Savanna ^c	Australian Forest ^d	North American Deciduous Forest ^e	Alaskan Boreal Forest ^f
	This Study	Hurst <i>et al.</i> [1994a] ^a	Hurst <i>et al.</i> , [1994b] ^a					
CO ₂	1613 ± 86	1595 ± 121	1646 ± 106	1722 ± 23	1640	1558	1671	1660
CO	88 ± 7	91	61	58	65	106	84.0	88.8
CH ₄	2.22 ± 0.32	2.34	2.36	1.31	2.4	3.60	5.18	2.79
C ₂ H ₆	0.53 ± 0.07	-	-	-	-	-	-	0.66
C ₂ H ₂	0.24 ± 0.03	0.06	0.14	-	-	-	-	0.24
C ₆ H ₆	0.21 ± 0.02	0.05	0.07	-	-	-	-	-

^aData originally presented in the different form [gC/gC DM (dry matter)] were converted into the common form [g species/kg DM].

^bFrom Ward *et al.* [1992].

^cFrom Andreae *et al.* [1996].

^dFrom Hurst *et al.* [1996].

^eFrom Yokelson *et al.* [1999a, 1999b].

^fFrom Goode *et al.* [2000].

reproducibility indicates that both data sets represent Australian savanna fires in the late dry season.

5. Emission Factors

[26] Emission factors for trace gases were derived from trace gas measurements in fresh smoke from fires. Commonly, an emission factor (EF) is defined as the mass of species released per unit mass of fuel consumed (g species/kg dry matter (DM)) [Ward *et al.*, 1992]. However, we start from a carbon mass balance method used by Hurst *et al.* [1994a, 1994b], which derives the fraction of the mass of carbon released to the atmosphere in the form of CO₂ relative to the dry mass of fuel carbon burned (g carbon/g carbon in dry matter (gC/gC DM)). The emission factor of CO₂ is calculated through the carbon balance equation shown below:

$$EF_{CO_2} = \frac{\Delta CO_2}{C_{fuel}} = \frac{\sum C_{emit}}{C_{fuel} \left(\frac{\Delta CO_2}{\Delta CO_2} + \frac{\Delta CO}{\Delta CO_2} + \frac{\Delta CH_4}{\Delta CO_2} + \frac{\Delta NMHC}{\Delta CO_2} + \frac{\Delta PC}{\Delta CO_2} \right)}$$

where PC = particulate carbon, $\sum C_{emit}$ = the mass of carbon released to the atmosphere, and C_{fuel} = the dry mass of fuel carbon burned.

[27] The fuel carbon content (46%), and the fraction of carbon emitted from the savanna fires to the atmosphere (96%) were taken from the weight and composition measurements of fuel and ash determined by Hurst *et al.* [1994a]. Fuels for the Hurst *et al.* [1994a] study were collected in the same northern Australian savanna in the late dry season as this study. A value of 0.088 ± 0.001 gC/gC DM was obtained for CO₂ and is in good agreement with the value of 0.087 ± 0.002 previously reported for Australian savanna fires by Hurst *et al.* [1994a, 1994b]. Emission factors of other compounds were determined using the same equation using the CO₂ fraction and emission ratios for the compounds of interest.

[28] Table 4 shows emission factors from this study converted to the common units of g species/kg DM [Ward *et al.*, 1992]. Comparing emission factors for different vegetation types, the variability for CO₂ was generally within the error range of different studies (Table 4). While the progress of the burning process for each ecosystem is largely characterized by its vegetation type, this result indicates that the emissions of CO₂ from a certain amounts

of biomass burned varies little according to the type of vegetation, after integration over the whole burning process for each ecosystem.

[29] By contrast, emission factors of CO and NMHCs vary significantly by ecosystem. This is closely related to the relationship between combustion efficiency and emission factors shown in Figure 4 (reproduced from Ward *et al.* [1992, Figure 8]). Strong linear correlations between combustion efficiency and emission factors of methane were shown based on combined data sets of previous field measurements which represent grassland (cerrado and savanna) and deforestation. Data from this study and Hurst *et al.* [1994a, 1994b] are plotted in Figure 4 as a solid triangle and a hollow triangle respectively, with the original two regression lines and the data points of Ward *et al.* [1992]. Our Australian data are reasonably distributed in the neighborhood of the model line for grassland ecosystems.

[30] These data show that more methane is emitted from forest fires than savanna fires of the same combustion efficiency. Higher methane emissions from forests are due to prolonged glowing combustion of logs, when alkanes and NH₃ are dominantly produced [Yokelson *et al.*, 1997]. The linear relationship between CO and other trace gases were presented in several other studies for both laboratory fires and the field fires [Hao *et al.*, 1996; Yokelson *et al.*, 1999a, 1999b]. Those results indicate that the variations of emission factors observed for different ecosystems can be explained by the different contribution ratios of each combustion process during which different gases are mainly produced. The proportion of each combustion process is mainly controlled by fuels and burning conditions of the ecosystems and emission factors of compounds mainly emitted during smoldering stage are expected to be greater for ecosystems with lower combustion efficiency. Because CO, CH₄, and most of the NMHCs are mainly emitted during the smoldering phase, the emission factors of those compounds were greater for forest fires than savanna fires (Table 4).

6. Estimation of Total Biomass Burned

[31] To estimate the gross emissions from biomass burning on a regional or global scale, we need to estimate the total biomass burned within the area of interest during the fire season. The estimation of total biomass burned is often accompanied by significant uncertainty, which can domi-




Figure 4. Methane production versus combustion efficiency shown for fires in grass-dominated and forested ecosystems (reproduced from *Ward et al.* [1992]).

nate the uncertainty of the gross emission estimates [*Seiler and Crutzen*, 1980; *Crutzen and Andreae*, 1990; *Hao and Liu*, 1994; *Scholes et al.*, 1996a, 1996b]. There have been two main methods (the “classification method” and the “modeling method”) for calculating annual emissions from tropical savanna burning. The classification method classifies the region of interest by vegetation types and extrapolate mean fuel load and area burned to each class. The modeling method sums the emissions from each fire, taking into account the actual fuel load at the time of the burning using a model [*Scholes et al.*, 1996a]. When fuel loads are relatively constant throughout the year, emissions can be estimated from the total areas of savanna and grassland burned annually. However in tropical savannas, there are large changes in the amount of grass fuels as the grasses develop over the growing season. Hence, it is very important to accurately follow the large fluctuations of fuel loads and area burned with a sufficient time resolution.

[32] In this study, fuel loads were estimated by a version of the modeling method improved by the Northern Territory Bushfire Council (NTBC), an agency which has the responsibility for fire prevention and management in rural areas of the Northern Territory, Australia [*Russell-Smith et al.*, 2003]. The NTBC calculation was based on estimates of the extent of burning derived from interpretation of satellite imagery and the quantity of fine fuels (<6 mm diameter) burned using empirical regional fuel accumulation relationships associated with detailed vegetation mapping. NOAA-AVHRR satellite images (pixels around 1.1 km²), were verified and calibrated by ground-verification studies and mapping of fire scars from fine-resolution LANDSAT TM

images (pixels 25 × 25 m). Derived values were 18.75 Mt DM yr⁻¹ for the region north of 15°S in the Northern Territory during the BIBLE-B period (2 weeks), and 212.3 Mt dm yr⁻¹ for the entire Australian savanna region in the year 1999. The confidence level of their fire extent mapping at larger regional scales was estimated to be 70–80%. The real amount of biomass burnt in 1999 cannot be less than the given estimates because heavier fuels (>6 mm diameter) were not accounted for in these estimates.

[33] A coarse estimation of the contribution of heavier fuels can be derived, however, from data presented by *Walker* [1981, Table 6]. Heavier fuel fractions (bark, shrubs, twigs) for Australian tropical savannas (*Walker's* region 3) are given as comprising typically 15% of total fuels (grass 65%, leaf litter 20%, bark 5%, shrubs 5%, and twigs 5%). *Walker's* tropical savanna region is, however, very extensive and most burning (including in 1999) occurred in higher rainfall regions of northern and northwestern Australia. Given evident relationships between both woody matter production and rainfall, and extent of late dry season (i.e., more intense) fires and consumption of woody fuels, it stands to reason that the proportion of heavier fuels consumed by fires in 1999 would be greater than the average 15% for the region as a whole. Conservatively, it is thus reasonable to assume that the heavier fuel component of savanna biomass burned in 1999 was at least 20% of all biomass burned in the tropical savanna region in 1999. Applying this qualification, we estimate that around 22 MT DM of biomass burned in the sample region during the 2 week period covering the BIBLE-B experiment, and around 255 MT DM was burned over the entire Australia savanna

Table 5. Source Strength Estimates for Australian Savanna Fires^a

	Emission Factor, g/kg dm	Emission Estimates, Gg/year
CH ₃ Br	0.0018 ± 0.0003	0.45 ± 0.29
CH ₃ Cl	0.0605 ± 0.0072	15 ± 10
CH ₃ I	0.0013 ± 0.0002	0.32 ± 0.2
C ₂ H ₆	0.13 ± 0.02	34 ± 21
C ₂ H ₄	0.31 ± 0.04	80 ± 50
C ₂ H ₂	0.12 ± 0.02	30 ± 19
C ₃ H ₈	0.027 ± 0.003	6.9 ± 4.2
C ₃ H ₆	0.071 ± 0.011	18 ± 12
i-C ₄ H ₁₀	0.0022 ± 0.0003	0.6 ± 0.4
C ₃ H ₄	0.0014 ± 0.0002	0.3 ± 0.2
n-C ₄ H ₁₀	0.0051 ± 0.0006	1.3 ± 0.8
t-2-C ₄ H ₈	0.0012 ± 0.0002	0.3 ± 0.2
1-C ₄ H ₈	0.010 ± 0.001	2.6 ± 1.7
c-2-C ₄ H ₈	0.00010 ± 0.0002	0.25 ± 0.17
i-C ₅ H ₁₂	0.0003 ± 0.0001	0.08 ± 0.05
n-C ₅ H ₁₂	0.0013 ± 0.0002	0.32 ± 0.20
C ₆ H ₁₄	0.0019 ± 0.0003	0.48 ± 0.31
C ₆ H ₆	0.035 ± 0.004	9.0 ± 5.6
C ₇ H ₁₆	0.00077 ± 0.00012	0.20 ± 0.13
Toluene	0.012 ± 0.001	3.0 ± 1.9
m-Xylene	0.0014 ± 0.0002	0.36 ± 0.23
p-Xylene	0.0004 ± 0.0001	0.10 ± 0.07
o-Xylene	0.0010 ± 0.0001	0.25 ± 0.16
CH ₄	2.2 ± 0.4	570 ± 380
CO	88 ± 8	23000 ± 13000
CO ₂	1613 ± 111	410000 ± 230000
DMS	0.0011	0.3

^aThese values are lower limits for reactive species. See text.

region in 1999. These values are employed in the following emission estimates.

7. Emission Estimates

[34] Table 5 shows our estimates of trace gas gross emissions based on the derived emission factors and total biomass burned. This is the first emission inventory for Australian savanna fires for most gases except CO₂, CO, CH₄ and a few NMHCs that were measured by *Hurst et al.* [1994a, 1994b]. For compounds with lifetimes longer than several hours, the relative uncertainty of the emission estimates in this study is estimated to be about +50%, close to the uncertainty estimates of *Scholes et al.* [1996a]. The estimated uncertainty is dominated by the substantial uncertainty in estimating biomass burnt (+50%) [*Russell-Smith et al.*, 2003], rather than other components such as uncertainty in the emission factors (±5%). For highly reactive compounds, the uncertainty in the emission factor increases substantially and could be even larger than the error in the

Table 6. Estimates of Total Biomass Burned and Emissions of CO₂ and CH₄ for Australian Savanna Fires

	CO ₂ , Tg C/year	CH ₄ , Tg C/year	Biomass Burned, Tg/year	CH ₄ /CO ₂
Current work	112 ± 64	0.42 ± 0.28	255 ± 128	0.0034
NGGIC [1998]	63	0.25	154	0.0042
Hurst et al. [1996]	69	0.28	173	0.0042
Hao et al. [1990]	174	-	425	-
Hao and Ward [1993]	-	0.30	290	0.0024




Figure 5. Area of land burned by prescribed burns of savanna and temperate grassland in Australia during 1990-1999 [NGGIC, 2000]. (NT: Northern Territory, Qld: Queensland, WA Western Australia).

estimation of biomass burned (i.e., ±75% for 1-butene). These errors were calculated using error propagation, considering each error associated with every parameter in the calculation (i.e., CO₂ gross emission: 410000 ± 230000 Gg ± 1613 ± 86 g/kg DM × 255 ± 128 Tg DM).

[35] The latest report on Australia's greenhouse gas emissions was released by the Australian *National Greenhouse Gas Inventory (NGGI)* [2000]. Ten-year averages of the emission estimates of CH₄, N₂O, NO_x, CO, and non-methane volatile organic compounds (NMVOC) from the prescribed burning of savanna and temperate grassland are given. The biomass burned was estimated by incorporating regional data for the biomass annually exposed to fire, the fraction of exposed biomass that burns, and the C content of the biomass. The average annual fuel loads and burning efficiencies (=0.72) were based on the review of *Tolhurst and Chatto* [1994], and the composition of the biomass and the emission factors were taken from *Hurst et al.* [1994a, 1994b].

[36] A tentative comparison of emission estimates from Australian savanna fires is shown in Table 6 [*Hao et al.*, 1990, 1996; *Hao and Ward*, 1993; *NGGI*, 2000; *Hurst et al.*, 1996; current study]. Our estimates of annual emission of CO₂ and CH₄ from savanna fires for 1999 are approximately 50% higher than the ten-year average presented by NGGIC. This difference is mainly due to higher fire activity in 1999 compared with the past ten years (Figure 5). This clearly suggests that our emission estimates are only for the specific year 1999 and cannot be applied for every year.

[37] Considering seasonal change of emission factors from biomass burning reported by *Hoffa et al.* [1999], emission factors are heavily affected by fuel moisture content and greater emission factors are expected in earlier dry season than later dry season for NMHCs. Thus the emission factors for NMHCs estimated in this study, based on the late dry season data might underestimate the emissions when applied to annual emission estimates. Accordingly, the effect of seasonal and annual variation of burning activity must be taken into account when using reported estimates of burning emissions.

8. Issues Involved in Emission Estimates

[38] To assess the contribution of biomass burning emissions to the global environment, a standardized data

set is of great importance. However, most of the emission estimates presented in previous studies were derived by various methods by various investigators [Seiler and Crutzen, 1980; Hao *et al.*, 1990; Hao and Ward, 1993; Delmas *et al.*, 1991]. In previous studies, Scholes *et al.* [1996a] pointed out that the amounts of biomass burned calculated by the “modeling method” were often substantially smaller than the amounts calculated by the “classification method”. Estimates of biomass burned using the classification method [Hao and Liu, 1994] were a factor of five higher than estimates obtained by the modeling method [Scholes *et al.*, 1996a]. This great discrepancy indicates the importance of knowing the method on which biomass burning estimates are based when comparing data.

[39] For instance, significant differences between estimates of the amount of biomass burned are apparent among the different studies in Table 6. The amount of total biomass burned presented by Hao *et al.* [1990] was first calculated by the classification method based on the data from Lacey *et al.* [1982] and then updated by Hao and Ward [1993] based on data from Galbally *et al.* [1992]. These estimates by Hao *et al.* were presented as mean annual estimates for the region and not corresponding to any particular year, while the rest of the studies were presented as estimates based on a single year. Meanwhile, the updated estimate of 290 Mt DM/yr [Hao and Ward, 1993] is lower than their previous value by approximately 30%, and is much closer to our estimate of 255 Mt DM/yr for 1999. Values of 154 Mt DM/yr for 1998 [NGGIC, 2000] and 173 Mt DM/yr for 1990 [Hurst *et al.*, 1996] were also derived by the classification method, but the tendency toward overestimation by the classification method noted previously for southern African fires by Scholes *et al.* [1996a] is not apparent, indicating the recent improvement of the classification method to prevent overestimating bias.

[40] In Table 6, the emission estimates for CO₂ directly reflect the variation of the estimated values of total biomass burned, but those for CH₄ show the effect of different emission ratios. The CH₄ emission factor used by Hao *et al.* [1996] was derived from the linear regression relationships as shown in Figure 4, based on data from various tropical regions while emission factors used in other studies shown in Table 6 were based on the field observation specifically investigated Australian savanna fires. The smaller value of CH₄ emission factors shown by Hao *et al.* [1996] is the result of a canceling effect of the estimates of the amount of biomass burned and emission factors as shown in Table 6. This illustrates the importance of recognizing combined effects of various factors when comparing emission estimates.

9. Future Studies

[41] As shown in the previous section where the variations among the estimates of the amount of biomass burned (Table 6) were much smaller compared to the discrepancy reported by Scholes *et al.* [1996a], convergence of the estimates by various methods is suggested for recent studies. Possible factors which contribute to reducing this discrepancy include the upgrade of inventories, observational results, and remote sensing data. Such improvements

might be made possible given substantial progress in field observational studies and remote sensing.

[42] Detailed studies on combustion mechanisms have provided us much useful information about trace gas emission from fires. Available data indicate that there are multiple factors that affect emissions such as burning efficiency (oxygen supply), burning temperature, fuel compositions, and vegetation characteristics, and it is necessary to clarify the relative importance of each factor and establish the relationship between emission strength and each factor. For this investigation, well-designed laboratory work and field studies are required.

[43] More remote sensing data are becoming available with the development of algorithms to quantitatively derive burning areas and burned biomass, using infrared or visible bands in most cases [Cahoon *et al.*, 1994; Menzel and Prins, 1996; Elvidge *et al.*, 1996; Setzer and Malingreau, 1996; Kendall *et al.*, 1997; Kauffman *et al.*, 1998]. Remote sensing is an efficient tool not only to obtain a precise estimate for certain areas and periods but to derive mean value of required parameters and to consider the magnitude of temporal variations once data are accumulated for enough long periods. As more efforts are made to improve algorithms and plans to launch sensors for fire detection unfold in the near future, utilization of remote sensing data will be one of the key factors to improve emission estimates.

[44] Tropospheric chemistry transport models (CTMs) are very efficient for considering the effect of trace gas emissions on the chemical composition and the oxidizing power of the global troposphere [Poisson *et al.*, 2000]. However, inventories provided for those models are not sufficient due to large uncertainties in emissions because some data are either not reported in national statistics and not reflecting fluctuations and trends during recent years [Olivier *et al.*, 1996]. For instance, Granier *et al.* [1999] found 25–50% differences in the amount of biomass burnt and CO₂ emitted when comparing results from their 3-D CTM and estimates by Andreae [1991]. These large discrepancies were considered to be mainly due to the scarcity of information on the amount of biomass burned in a given year and the large year-to-year variations in the amount of biomass burned. Data sets with higher spatial and temporal resolution are greatly needed as input for models.

[45] Finally, to build a reliable data set, available data must be standardized and integrated. The approaches for this integration are getting started recently [Andreae and Merlet, 2001]. The reasonable assessment of the effect of biomass burning emission to the global environment will be possible when all the above listed progress were achieved in the near future.

10. Conclusions

[46] Selected atmospheric trace gases emitted by Australian savanna fires were measured in the local late dry season. Emission ratios and emission factors were determined for various trace gases based on BL enhancements in the source region. The emission ratios obtained for Australian savanna fires agreed well with previous measurements for Brazil and Africa for stable compounds such as CH₄ and alkanes, while lower ratios were observed for

highly reactive compounds such as alkenes affected by oxidation during transport. The emission factors depended heavily on burning conditions such as combustion efficiency. Emission factors of CO, CH₄ and NMHCs varied significantly among different vegetation types, while those of CO₂ varied little.

[47] The annual trace gas emissions of Australian savanna fires were deduced from the emission factors and estimates of burned fuel using a modeling method appropriate for 1999. Emission parameters such as emission ratios and emission factors were based on an excellent data set obtained by aircraft observations and appropriately interpreted remote sensing data. As a result, uncertainty in estimates of Australian savanna fire emissions was considerably reduced compared to previous studies. While significant uncertainty still remains in estimates of the total biomass burned, mainly due to the difficulty of taking spatial and temporal variation into account, diminishing discrepancies among different studies indicate a general improvement in methods and databases. As shown in this study, the use of remote sensing data has a great potential to provide databases with sufficient spatial/temporal resolution. Approaches for integrating component data sets are now being developed which, in future years, will facilitate more reliable estimates of emissions from biomass burning.

[48] **Acknowledgments.** The authors gratefully acknowledge the members of the BIBLE science team. Special thanks to the pilots and the crew of the Diamond Air Service for performing the aircraft operations. Thanks also to the staff of the Pel-Air Aviation and those of the Northern Territory Bushfire Council for their excellent support during the campaign in Darwin. We particularly would like to thank Murray McEachern, Ben Liley and Brent Love for kind assistance with sampling, and the team at UCI for their excellent laboratory work and data processing to produce this NMHC and halocarbon data set. We acknowledge the valuable comments and suggestions of Robert Yokelson, Nicola Blake, David Streets, Isobel Simpson and Yugo Kanaya. The BIBLE-B mission was carried out under the research program of Earth Observation Research Center, National Space Development Agency of Japan (NASDA/EORC).

References

- Andreae, M. O., Biomass burning: Its history, use, and distribution and its impact on environmental quality and global climate, in *Global Biomass Burning: Atmospheric, Climatic, and Biospheric Implications*, edited by J. S. Levine, pp. 3–21, MIT Press, Cambridge, Mass., 1991.
- Andreae, M. O., and P. Merlet, Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cycles*, **15**, 955–966, 2001.
- Andreae, M. O., B. E. Anderson, D. R. Blake, J. D. Bradshaw, J. E. Collins, G. L. Gregory, G. W. Sachse, and M. C. Shipham, Influence of plumes from biomass burning on atmospheric chemistry over the equatorial Atlantic during CITE-3, *J. Geophys. Res.*, **99**, 12,793–12,808, 1994.
- Andreae, M. O., et al., Methyl halide emission from savanna fires in southern Africa, *J. Geophys. Res.*, **101**, 23,603–23,614, 1996.
- Blake, D. R., D. F. Hurst, T. W. Smith Jr., W. J. Whipple, T.-Y. Chen, N. J. Blake, and F. S. Rowland, Summertime measurements of selected nonmethane hydrocarbons in the Arctic and sub-Arctic during the 1988 Arctic Boundary Layer Expedition (ABLE 3A), *J. Geophys. Res.*, **97**, 16,559–16,588, 1992.
- Blake, D. R., T. W. Smith Jr., T.-Y. Chen, W. J. Whipple, and F. S. Rowland, Effect of biomass burning on summertime nonmethane hydrocarbons concentrations in the Canadian wetlands, *J. Geophys. Res.*, **99**, 1699–1719, 1994.
- Blake, N. J., D. R. Blake, B. C. Sive, T.-Y. Chen, F. S. Rowland, J. E. Collins Jr., G. W. Sachse, and B. E. Anderson, Biomass burning emissions and vertical distribution of atmospheric methyl halides and other reduced carbon gases in the South Atlantic region, *J. Geophys. Res.*, **101**, 24,151–24,164, 1996a.
- Blake, N. J., D. R. Blake, J. E. Collins Jr., G. W. Sachse, B. E. Anderson, J. A. Brass, P. J. Riggan, and F. S. Rowland, Biomass burning emissions of atmospheric methyl halide and hydrocarbon gases in the South Atlantic region, in *Biomass Burning and Global Change*, edited by J. S. Levine, pp. 575–594, MIT Press, Cambridge, Mass., 1996b.
- Blake, N. J., et al., Influence of southern hemispheric biomass burning on mid-tropospheric distributions of nonmethane hydrocarbons and selected halocarbons over the remote South Pacific, *J. Geophys. Res.*, **104**, 16,213–16,232, 1999.
- Bonsang, B., G. Lambert, and C. C. Boissard, Light hydrocarbons emissions from African savanna burnings, in *Global Biomass Burning: Atmospheric, Climatic, and Biospheric Implications*, edited by J. S. Levine, pp. 155–161, MIT Press, Cambridge, Mass., 1991.
- Browell, E. V., et al., Air mass characteristics observed over the tropical south Atlantic, Africa, and Brazil during the springtime, *J. Geophys. Res.*, **101**, 24,043–24,068, 1996.
- Butler, J. H., Atmospheric chemistry: Better budgets for methyl halides?, *Nature*, **403**, 260–261, 2000.
- Cahoon, D. R., Jr., B. J. Stocks, J. S. Levine, W. R. Cofer III, and J. M. Pierson, Satellite analysis of the severe 1987 forest fires in northern China and southeastern Siberia, *J. Geophys. Res.*, **99**, 18,627–18,638, 1994.
- Charlson, R. J., J. E. Lovelock, M. O. Andreae, and S. G. Warren, Oceanic phytoplankton, atmospheric sulphur, cloud albedo and climate, *Nature*, **326**, 655–661, 1987.
- Colman, J. J., A. L. Swanson, S. Meinardi, B. C. Sive, D. R. Blake, and F. S. Rowland, Description of the analysis of a wide range of volatile organic compounds in whole air samples collected during PEM-Tropics A and B, *Anal. Chem.*, **73**, 3723–3731, 2001.
- Connors, V. S., M. Flood, T. Jones, B. Gormsen, S. Nolf, and H. G. Reichle Jr., Global distribution of biomass burning and carbon monoxide in the middle troposphere during early April and October 1994, in *Biomass Burning and Global Change*, edited by J. S. Levine, pp. 99–106, MIT Press, Cambridge, Mass., 1996.
- Crutzen, P. J., and M. O. Andreae, Biomass burning in the tropics: Impact on atmospheric chemistry and biogeochemical cycles, *Science*, **250**, 1669–1698, 1990.
- Crutzen, P. J., A. C. Delany, J. Greenberg, P. Haagenson, L. Heidt, R. Lueb, W. Pollock, W. Seiler, A. Wartburg, and P. Zimmerman, Tropospheric chemical composition measurements in Brazil during the dry season, *J. Atmos. Chem.*, **2**, 233–256, 1985.
- Delmas, R. A., On the emission of carbon, nitrogen, and sulphur in the atmosphere during bush fires in intertropical savanna zones, *Geophys. Res. Lett.*, **9**, 761–764, 1982.
- Elvidge, C. D., H. W. Kroehl, E. A. Kihn, K. E. Baugh, E. R. Davis, and W. M. Hao, Algorithm for the retrieval of fire pixels from DMSP Operational Linescan System, in *Biomass Burning and Global Change*, edited by J. S. Levine, pp. 73–85, MIT Press, Cambridge, Mass., 1996.
- Ferek, R. J., J. S. Reid, P. V. Hobbs, D. R. Blake, and C. Lioussis, Emission factors of hydrocarbons, halocarbons, trace gases and particles from biomass burning in Brazil, *J. Geophys. Res.*, **103**, 32,107–32,118, 1998.
- Folkens, I. A., A. J. Weinheimer, B. A. Ridley, J. G. Walega, B. Anderson, J. E. Collins, G. Sachse, R. F. Pueschel, and D. R. Blake, O₃, NO_y, and NO_x/NO_y in the upper troposphere of the equatorial Pacific, *J. Geophys. Res.*, **100**, 20,913–20,926, 1995.
- Galbally, I. E., P. J. Fraser, C. P. Meyer, and D. W. T. Griffith, Biosphere-atmosphere exchange of trace gases over Australia, in *Australia's Renewable Resources: Sustainability and Global Change*, *BRR Proc. 14*, edited by R. M. Gifford and M. M. Barson, pp. 118–149, Bur. of Rural Resour., Commonw. Sci. and Ind. Res. Org., Div. of Plant Indust., Melbourne, Vict., Australia, 1992.
- Goode, J. G., R. J. Yokelson, D. E. Ward, R. A. Susott, R. E. Babbitt, M. A. Davies, and W. M. Hao, Measurements of excess O₃, CO₂, CO, CH₄, C₂H₄, C₂H₂, HCN, NO, NH₃, HCOOH, CH₃COOH, HCHO, and CH₃OH in 1997 Alaskan biomass burning plumes by airborne Fourier transform infrared spectroscopy (AFTIR), *J. Geophys. Res.*, **105**, 22,147–22,166, 2000.
- Granier, C., J. F. Muller, and G. Brasseur, The impact of biomass burning on the global budget of ozone and ozone precursors, in *Proceedings of the Wengen Conference on Global Change Research: Biomass Burning and its Inter-Relationships with the Climate System*, pp. 69–84, 1999.
- Greenberg, J. P., P. R. Zimmerman, L. Heidt, and W. Pollock, Hydrocarbon and carbon monoxide emissions from biomass burning in Brazil, *J. Geophys. Res.*, **89**, 1350–1354, 1984.
- Gregory, G. L., H. E. Fuelberg, S. P. Longmore, B. E. Anderson, J. E. Collins, and D. R. Blake, Chemical characteristics of tropospheric air over the tropical South Atlantic Ocean: Relationship to trajectory history, *J. Geophys. Res.*, **101**, 23,957–23,972, 1996.
- Gregory, G. L., et al., Chemical characteristics of Pacific tropospheric air in the region of the Intertropical Convergence Zone and South Pacific Convergence Zone, *J. Geophys. Res.*, **104**, 5677–5696, 1999.
- Hao, W. M., and D. E. Ward, Methane production from global biomass burning, *J. Geophys. Res.*, **98**, 20,657–20,661, 1993.

- Hao, W. M., and M. H. Liu, Spatial and temporal distribution of tropical biomass burning, *Global Biogeochem. Cycles*, **8**, 495–503, 1994.
- Hao, W. M., M. H. Liu, and P. J. Crutzen, Estimates of annual and regional releases of CO₂ and other trace gases to the atmosphere from fires in the tropics, based on the FAO statistics for the period 1975–1980, in *Fire in the Tropical Biota: Ecosystem Processes and Global Challenges*, edited by J. G. Goldammer, pp. 440–462, Springer-Verlag, New York, 1990.
- Hao, W. M., D. E. Ward, G. Olbu, and S. P. Baker, Emissions of CO₂, CO, and hydrocarbons from fires in diverse African savanna ecosystems, *J. Geophys. Res.*, **101**, 23,577–23,584, 1996.
- Hoffa, E. A., D. E. Ward, W. M. Hao, R. A. Susott, and R. H. Wakimoto, Seasonality of carbon emissions from biomass burning in a Zambian savanna, *J. Geophys. Res.*, **104**, 13,841–13,853, 1999.
- Hurst, D. F., D. W. T. Griffith, and G. D. Cook, Trace gas emissions and biomass burning in tropical Australian savannas, *J. Geophys. Res.*, **99**, 16,441–16,456, 1994a.
- Hurst, D. F., D. W. T. Griffith, J. N. Carras, D. J. Williams, and P. J. Fraser, Measurements of trace gas emitted by Australian savanna fires during the 1990 dry season, *J. Atmos. Chem.*, **18**, 33–56, 1994b.
- Hurst, D. F., D. W. T. Griffith, and G. D. Cook, Trace gas emissions from biomass burning in Australia, in *Biomass Burning and Global Change*, edited by J. S. Levine, pp. 788–792, MIT Press, Cambridge, Mass., 1996.
- Jonquière, I., A. Marengo, A. Maalej, and F. Rohrer, Study of ozone formation and transatlantic transport from biomass burning emissions over West Africa during the airborne Tropospheric Ozone Campaigns TROPOZ I and TROPOZ II, *J. Geophys. Res.*, **103**, 19,059–19,073, 1998.
- Kauffman, J. B., D. L. Cummings, and D. E. Ward, Relationships of fire, biomass and nutrient dynamics along a vegetation gradient in the Brazilian cerrado, *J. Ecol.*, **82**, 519–531, 1994.
- Kauffman, J. B., C. O. Justice, L. P. Flynn, J. D. Kendall, E. M. Prins, L. Gigilio, D. E. Ward, W. P. Menzel, and A. W. Setzer, Potential global fire monitoring from EOS-MODIS, *J. Geophys. Res.*, **103**, 32,215–32,238, 1998.
- Kendall, J. D., C. O. Justice, P. R. Dowty, C. D. Elvidge, and J. G. Goldammer, Remote sensing of fires in southern Africa during the SAFARI 1992 campaign, in *Fire in Southern African Savannas: Ecological and Atmospheric Perspectives*, edited by B. W. van Wilgen et al., pp. 89–133, Witwatersrand Univ. Press, Johannesburg, 1997.
- Kondo, Y., et al., Effects of biomass burning, lightning, and convection on O₃, CO, and NO_y over the tropical Pacific and Australia in August–October 1998 and 1999, *J. Geophys. Res.*, **107**, 8402, doi:10.1029/2001JD000820, 2002 [printed 108(D3)], 2003].
- Lacey, C. J., J. Walker, and I. R. Noble, Fire in Australian tropical savannas, in *Ecology of Tropical Savannas*, edited by B. J. Huntley and B. H. Walker, pp. 246–272, Springer-Verlag, New York, 1982.
- Lobert, J. M., D. H. Scharffe, W. M. Hao, T. A. Kuhlbusch, R. Seuwen, P. Warneck, and P. J. Crutzen, Experimental evaluation of biomass burning emissions: Nitrogen and carbon containing compounds, in *Global Biomass Burning: Atmospheric, Climatic, and Biospheric Implications*, edited by J. S. Levine, pp. 289–304, MIT Press, Cambridge, Mass., 1991.
- Machida, T., K. Kita, Y. Kondo, D. Blake, S. Kawakami, G. Inoue, and T. Ogawa, Vertical and meridional distributions of the atmospheric CO₂ mixing ratio between northern midlatitudes and southern subtropics, *J. Geophys. Res.*, **8401**, doi:10.1029/2001JD000910, 2002 [printed 108(D3)], 2003].
- Mauzerall, D. L., J. Logan, D. J. Jacob, B. E. Anderson, D. R. Blake, J. D. Bradshaw, B. Heikes, G. W. Sachse, H. Singh, and B. Talbot, Seasonal characteristics of tropospheric ozone production and mixing ratios over East Asia: A global three-dimensional chemical transport model analysis, *J. Geophys. Res.*, **103**, 8401–8423, 1998.
- McKenzie, L. M., D. E. Ward, and W. M. Hao, Chlorine and bromine in the biomass of tropical and temperate ecosystems, in *Biomass Burning and Global Change*, edited by J. S. Levine, pp. 241–248, MIT Press, Cambridge, Mass., 1996.
- Menzel, W. P., and E. M. Prins, Monitoring biomass burning with the new generation of geostationary satellites, in *Biomass Burning and Global Change*, edited by J. S. Levine, pp. 56–64, MIT Press, Cambridge, Mass., 1996.
- National Greenhouse Gas Inventory (NGGIC), *National Greenhouse Gas Inventory 1998*, Nat. Greenhouse Gas Invent. Comm., Canberra, Australia, 2000.
- Olivier, J. G. J., A. F. Bouwman, C. W. M. van der Maas, J. J. M. Berdowski, C. Veldt, J. P. J. Bloos, A. J. H. Visschedijk, Y. J. Zandveld, and J. L. Haverlag, Description of EDGAR version 2.0: A set of global emission inventories of greenhouse gases and ozone-depleting substances for all anthropogenic and most natural sources on a per country basis and on a 1 × 1 degree grid, *RIVM Rep. 771060 002/TNO-MEP Rep. R96/119*, Nat. Inst. of Public Health and the Environ., Bilthoven, Netherlands, 1996.
- Poisson, N., M. Kanakidou, and P. J. Crutzen, Impact of non-methane hydrocarbons on tropospheric chemistry and the oxidizing power of the global troposphere: 3-dimensional modelling results, *J. Atmos. Chem.*, **36**, 157–230, 2000.
- Reinhardt, T. E., and D. E. Ward, Factors affecting methyl chloride emissions from forest biomass combustion, *Environ. Sci. Technol.*, **29**, 825–832, 1995.
- Ridley and Atlas, Role of active nitrogen in odd-hydrogen partitioning, in *Atmospheric chemistry and global change*, edited by G. P. Brasseur, J. J. Orlando, and G. S. Tyndall, pp. 268–275, Oxford Univ. Press, New York, 1999.
- Russell-Smith, J., A. C. Edwards, and G. D. Cook, Reliability of biomass burning estimates from savanna fires: Biomass burning in northern Australia during the 1999 Biomass Burning and Lightning Experiment B field campaign, *J. Geophys. Res.*, **108**, doi:10.1029/2001JD000787, in press, 2003.
- Scholes, R. J., J. Kendall, and C. O. Justice, The quantity of biomass burned in southern Africa, *J. Geophys. Res.*, **101**, 23,667–23,676, 1996a.
- Scholes, R. J., D. E. Ward, and C. O. Justice, Emissions of trace gases and aerosol particles due to vegetation burning in southern hemisphere Africa, *J. Geophys. Res.*, **101**, 23,677–23,682, 1996b.
- Seiler, W., and P. J. Crutzen, Estimates of gross and net fluxes of carbon between the biosphere and atmosphere from biomass burning, *Clim. Change*, **2**, 247–297, 1980.
- Setzer, A. W., and J. P. Malingreau, AVHRR monitoring of vegetation fires in the tropics: Towards a global product, in *Biomass Burning and Global Change*, edited by J. S. Levine, pp. 48–81, MIT Press, Cambridge, Mass., 1996.
- Singh, H. B., et al., Biomass burning influences on the composition of the remote South Pacific troposphere: Analysis based on observations from PEM-Tropics-A, *Atmos. Environ.*, **34**, 635–644, 2000.
- Sive, B. C., Atmospheric NMHCs: Analytical methods and estimated hydroxyl radical concentrations, Ph.D. dissertation, Univ. of Calif., Irvine, 1998.
- Takegawa, N., et al., Airborne vacuum ultraviolet resonance fluorescence instrument for in situ measurement of CO, *J. Geophys. Res.*, **106**, 24,237–24,244, 2001.
- Takegawa, N., et al., Removal of NO_x and NO_y in biomass burning plumes in the boundary layer over northern Australia, *J. Geophys. Res.*, **108**, doi:10.1029/2002JD002505, in press, 2003.
- Talbot, R. W., et al., Chemical characteristics of continental outflow over the tropical south Atlantic Ocean from Brazil and Africa, *J. Geophys. Res.*, **101**, 24,187–24,202, 1996.
- Tolhurst, K. G., and K. Chatto, Inventory of areas burnt and fuels consumed by bushfires in Australia 1983 to 1994, *Rep. for Commonw. Sci. and Ind. Res. Org., Div. of Atmos. Res.*, **17** pp., Cent. for Forest Tree Technol., 1994.
- Walker, J., Fuel dynamics in Australian vegetation, in *Fire and the Australian Biota*, edited by A. M. Gill, R. M. Groves, and I. R. Noble, pp. 101–127, Austr. Acad. of Sci., Canberra, 1981.
- Ward, D. E., and L. F. Radke, Emission measurements from vegetation fires: A comparative evaluation of methods and results, in *Fire in the Environment: The Ecological, Atmospheric and Climatic Importance of Vegetation Fires*, edited by P. J. Crutzen and J. G. Goldammer, pp. 53–76, John Wiley, New York, 1993.
- Ward, D. E., R. A. Susott, J. B. Kauffman, R. E. Babbitt, D. L. Cummings, B. Dias, B. N. Holben, Y. J. Kaufman, R. A. Rasmussen, and A. W. Setzer, Smoke and fire characteristics for cerrado and deforestation burns in Brazil: BASE B experiment, *J. Geophys. Res.*, **97**, 14,601–14,619, 1992.
- Warneck, P., *Chemistry of the Natural Atmosphere*, edited by R. Dmowska, J. R. Holton, and H. T. Rossby, pp. 594–611, Academic, San Diego, Calif., 2000.
- World Meteorological Organization (WMO), *Scientific Assessment of Ozone Depletion: 1998*, Global Ozone Res. and Monit. Proj., Geneva, 1999.
- Yokelson, R. J., D. W. Griffith, and D. E. Ward, Open-path Fourier transform infrared studies of large-scale laboratory biomass fires, *J. Geophys. Res.*, **101**, 21,067–21,080, 1996.
- Yokelson, R. J., R. A. Susott, D. E. Ward, J. Readon, and D. W. Griffith, Emissions from smoldering combustion of biomass measured by open-path Fourier transform infrared spectroscopy, *J. Geophys. Res.*, **102**, 18,865–18,877, 1997.
- Yokelson, R. J., J. G. Goode, D. E. Ward, R. A. Susott, R. E. Babbitt, D. D. Wade, I. Bertschi, D. W. T. Griffith, and W. M. Hao, Emissions of formaldehyde, acetic acid, methanol, and other trace gases from biomass fires in North Carolina measured by airborne Fourier transform infrared spectroscopy, *J. Geophys. Res.*, **104**, 30,109–30,125, 1999a.

- Yokelson, R. J., R. A. Susott, R. E. Babbitt, W. M. Hao, and D. E. Ward, Trace gas emissions from specific biomass fire-types (abstract), in *START Synthesis Workshop on Greenhouse Gas Emission, Aerosols and Land Use and Cover Change in Southeast Asia*, edited by T. Moya, pp. 60–68, SARCS, IGAC, LUCC, IGBP China-Taipei, Taipei, 1999b.
-
- D. R. Blake, S. Meinardi, and F. S. Rowland, Department of Chemistry, 570 Rowland Hall, University of California-Irvine, Irvine, CA 92697-2025, USA.
- A. Edwards and J. Russell-Smith, Tropical Savannas Cooperative Research Centre, Bushfires Council of the Northern Territory, P. O. Box 37346, Winnellie 0821, Northern Territory, Australia.
- S. Kawakami, T. Ogawa, and T. Shirai, National Space Development Agency of Japan, Earth Observation Research Center, 23F Office Tower X, Triton Square, 1-8-10 Harumi, Chuo-ku, Tokyo 104-6023, Japan. (shirai@uci.edu)
- K. Kita, Department of Environmental Sciences Faculty of Science, Ibaraki University, 2-1-1 Bunkyo, Mito, Ibaraki 310-8512, Japan.
- Y. Kondo, Research Center for Advanced Science and Technology, University of Tokyo, 4-6-1 Komaba, Meguro-ku, Tokyo, 153-8904, Japan.
- M. Koike, Department of Earth and Planetary Sciences, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan.
- T. Machida, National Institute for Environmental Studies, 16-2 Onogawa, Tsukuba, 305-8509, Japan.
- N. Nishi, Department of Earth and Planetary Science, Kyoto University, Kitashirakawa-Oiwake-cho, Sakyo-ku, Kyoto, 606-8502, Japan.
- N. Takegawa, Solar-Terrestrial Environment Laboratory, Nagoya University, 3-13, Honohara, Toyokawa, Aichi, 442-8507 Japan.